

# REGENESIS VET™ R.I.A.D.™ DOSING & ADMINISTRATION

## RECOMMENDED DOSING

- Each Regenes Vet™ R.I.A.D.™ treatment should be 15-20ml in volume
- Normal sterility standards should be adhered to when administering Regenes Vet™ R.I.A.D.™
- Please familiarize yourself with the full list of warnings and contraindications in the prescribing information before proceeding with treatment
- Intravenous administration
  - Regenes Vet™ R.I.A.D.™ should be administered **very slowly and evenly** at a rate no faster than 1ml/4 seconds via intravenous route of administration
  - PLEASE NOTE: Faster injection rates increase the risk of an adverse event
- Intramuscular administration
  - Regenes Vet™ R.I.A.D.™ should be administered **very slowly and evenly** at a rate no faster than 1ml/2 seconds via intramuscular route of administration
  - The full treatment can be split evenly and injected at 2 different injection sites
- Treatments should be frozen within 2-3 hours of processing
- Treatments can be stored for up to 24 months provided the cold chain is not interrupted (kept frozen)
- Cold chain interruptions could adversely affect the quality of the product and the sterility
- Defrost at room temperature out of direct sunlight one hour before administration
- **DO NOT** ADMINISTER REGENESIS VET™ R.I.A.D.™ WITHIN 21 DAYS OF AN AFRICAN HORSE SICKNESS (AHS) VACCINE OR WITHIN 14 DAYS OF ANY OTHER VACCINE
- TO BE ADMINISTERED BY A LICENSED VETERINARIAN **ONLY**

## RECOMMENDED FREQUENCY

### MAINTENANCE

1 x injection on a monthly basis



### INJURY

1 x injection every 10 days for  
3 injections



### PRE-SHOW/TRAVEL

7-10 days before show or travel



### AFRICAN HORSE SICKNESS VACCINE

